

Quantifying Methane, Ethane, and VOC Emissions from the Denver-Julesburg Basin Using Aircraft Mass Balance in Fall of 2021.


Hannah M. Daley¹, Alan Fried², Xinrong Ren³, Phillip Stratton³, Benjamin Hmiel⁴ and Russell R. Dickerson¹.

¹Department of Atmospheric and Oceanic Science, University of Maryland; ²University of Colorado at Boulder, Institute for Arctic and Alpine Research, Boulder, United States; ³Air Resources Laboratory, National Oceanic and Atmospheric Administration; ⁴Colorado Department of Public Health and Environment, Denver, United States.

Intro:

Methane, a potent greenhouse gas and primary component of natural gas, requires quantification of emission rates and their temporal changes to assess its climatic impact. In the Denver-Julesburg Basin (DJB), particularly the Wattenberg field area of Colorado, methane emissions predominantly originate from oil and natural gas (O&G) operations (>18,000 active wells in 2021), agricultural activities at concentrated animal feedlot operations (CAFOs), and landfills (Figure 2). During Fall 2021 (9/17–10/5), our research group conducted nine flights over this region in a fully instrumented Cessna aircraft (Figure 1), measuring CH₄, C₂H₆, acetic acid, and other trace gases, with four flights deemed sufficient for mass balance emission rate determination. Since CH₄ and C₂H₆ are co-emitted from O&G operations, and CH₄ and acetic acid (but not C₂H₆) from CAFOs (Figure 3), we determined methane emission rates, source apportionment, and emission intensity for each flight normalized by O&G production (Figure 4).

Aircraft-Based Mass Balance Method:



$$CH_4 ER = \int_{z_0=surface; y_0}^{z_1; y_1} ([CH_4]_{plume} - [CH_4]_{bkg}) * WS_{\perp} * (GS * \Delta t) * N(P, T, Z) dy dz$$

$$WS_{\perp} = WS * \cos(WD_{\perp} - TA) * k$$

Emission uncertainties were assessed through a propagation error analysis of 10 variables: adjusted PBLH, methane concentration ($[CH_4]_{plume}$), methane background ($[CH_4]_{bkg}$), wind speed (WS), track angle (TA), wind speed correction factor (k), wind direction (WD), groundspeed (GS), temperature (T), and pressure (P).

Emission Intensity Determination:

$$Emission Intensity (EI) = (CH_4 \text{ fraction from O\&G}) * (CH_4 \text{ mass balance kg/hour}) / (BOEh)$$

Methane and Ethane Emission Results:

Date	C2H6 Emission Rate (tn/hr)	CH4 Emission Rate (tn/hr)	O&G%	Total BOE/hr	Methane Emission Intensity	Ethane Emission Intensity
9/24/2021	2.24 ± 0.49	17.7 ± 4.3	51.4	36925	0.26	0.06 ± 0.1
9/27/2021	1.83 ± 0.50	17.9 ± 2.9	63.2	27951	0.40	0.07 ± 0.02
10/1/2021	1.96 ± 0.52	19.5 ± 7.3	71.2	17949	0.77	0.11 ± 0.04
10/5/2021	3.71 ± 0.59	28.5 ± 3.1	67.1	29337	0.65	0.13 ± 0.02
Average	2.44 ± 0.80	20.9 ± 5.0	63.2 ± 8.7	28041 ± 7800	0.56 ± 0.19	0.09 ± 0.02

As oil and gas production in the D-J Basin increases, methane emission intensity decreases.



Poster number: A21K – 1917



Scan for a link to the poster, reference information, and more information about the presenter.

3 Take away points:

- 1. Methane and Ethane Emissions:** From four flights conducted in Fall 2021 (9/24–10/5), the methane emission rate was quantified as 20.9 ± 5.9 tons CH₄/hr, while the ethane emission rate was 2.44 ± 0.80 tons C₂H₆/hr.
- 2. Source Apportionment Using Tracers:** Acetic acid proved to be a valuable tracer for agricultural methane emissions. Combining acetic acid and ethane tracers, we attributed $63.2 \pm 8.7\%$ of methane emissions to oil and gas (O&G) operations.
- 3. Emission Intensity Trends:** By determining flight footprints, we achieved a more precise estimation of O&G production captured during each leg/flight. This enabled calculations of a methane emission intensity of 0.56 ± 0.19 kg thermogenic CH₄/BOE/hr and an ethane emission intensity of 0.09 ± 0.02 kg C₂H₆/BOE/hr. Methane emission intensity has declined linearly by 0.05 kg/BOE/hr per year since 2012, while ethane emission intensity has significantly decreased > 70% since 2015 (Pieschl, 2018; Ngulat/Daley/Fried et al., in prep.).




Figure 1. (Top left) The 402B aircraft and reverse-facing gas inlet, along with pressure/temperature/humidity sensors. (Top middle) Interior view showing the UMD instrument rack and CAMS-2 ethane instrument. (Top right) The PTR-TOF spectrometer inside the aircraft. (Bottom) Schematic of instrument placements relative to the aircraft size.




Figure 2. Map of the Wattenberg Field portion of the DJB in Weld and Larimer Counties showing CH₄ sources, with O&G facilities sized by daily BOE production and CAFOs sized by animal units (AU).

Acetic Acid: A Key Tracer for CAFO Emissions




Figure 3. Plot of 1s ΔMethane vs. ΔEthane, colored and sized by Δacetic acid concentrations. Background values represent 5% of values within the PBLH. The box region highlights Greeley, CO, a major CAFO area. The figure shows a ΔEthane/ΔMethane ratio of approximately 5.5% and a strong correlation between CAFOs and acetic acid.




Figure 4. Methane emission intensity (blue) and Barrels of Oil Equivalent (BOE, black) for six studies conducted in the Denver-Julesburg Basin from 2012–2021. BOE increased by 146% from 2012 to 2021 and 137% from 2015 to 2021. The trendline ($r^2 = 0.78$) indicates a decrease in emission intensity of approximately 0.05 EI/year, where EI is kg CH₄/BOE/hr.