

# Satellite retrieval of cloud condensation nuclei concentrations by using clouds as CCN chambers

Daniel Rosenfeld<sup>a,1</sup>, Youtong Zheng<sup>b,c,d</sup>, Eyal Hashimshoni<sup>a</sup>, Mira L. Pöhlker<sup>e,f</sup>, Anne Jefferson<sup>g</sup>, Christopher Pöhlker<sup>e</sup>, Xing Yu<sup>h</sup>, Yannian Zhu<sup>d,h</sup>, Guihua Liu<sup>h</sup>, Zhiguo Yue<sup>h</sup>, Baruch Fischman<sup>a</sup>, Zhanqing Li<sup>b,c,d</sup>, David Giguzin<sup>a</sup>, Tom Goren<sup>a</sup>, Paulo Artaxo<sup>i</sup>, Henrique M. J. Barbosa<sup>i</sup>, Ulrich Pöschl<sup>e,f</sup>, and Meinrat O. Andreae<sup>e</sup>

<sup>a</sup>Institute of Earth Sciences, The Hebrew University of Jerusalem, Jerusalem 91904, Israel; <sup>b</sup>Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD 20740; <sup>c</sup>Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD 20740; <sup>d</sup>State Key Laboratory of Earth Surface Processes and Resource Ecology, College of Global Change and Earth System Science, Beijing Normal University, Beijing 100875, China; <sup>e</sup>Biogeochemistry Department, Max Planck Institute for Chemistry, Mainz 55020, Germany; <sup>f</sup>Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz 55020, Germany; <sup>g</sup>Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO 80305; <sup>h</sup>Meteorological Institute of Shaanxi Province, Xi'an 710015, China; and <sup>i</sup>Physics Institute, University of Sao Paulo, Sao Paulo 05508-090, Brazil

Edited by John H. Seinfeld, California Institute of Technology, Pasadena, CA, and approved January 29, 2016 (received for review September 14, 2015)

**Quantifying the aerosol/cloud-mediated radiative effect at a global scale requires simultaneous satellite retrievals of cloud condensation nuclei (CCN) concentrations and cloud base updraft velocities ( $W_b$ ). Hitherto, the inability to do so has been a major cause of high uncertainty regarding anthropogenic aerosol/cloud-mediated radiative forcing. This can be addressed by the emerging capability of estimating CCN and  $W_b$  of boundary layer convective clouds from an operational polar orbiting weather satellite. Our methodology uses such clouds as an effective analog for CCN chambers. The cloud base supersaturation ( $S$ ) is determined by  $W_b$  and the satellite-retrieved cloud base drop concentrations ( $N_{db}$ ), which is the same as  $CCN(S)$ . Validation against ground-based CCN instruments at Oklahoma, at Manaus, and onboard a ship in the northeast Pacific showed a retrieval accuracy of  $\pm 25\%$  to  $\pm 30\%$  for individual satellite overpasses. The methodology is presently limited to boundary layer not raining convective clouds of at least 1 km depth that are not obscured by upper layer clouds, including semi-transparent cirrus. The limitation for small solar backscattering angles of  $< 25^\circ$  restricts the satellite coverage to  $\sim 25\%$  of the world area in a single day.**

CCN concentrations | satellite remote sensing | convective clouds | cloud–aerosol interactions

The Intergovernmental Panel on Climate Change (IPCC) report (1) states that the uncertainty in aerosol/cloud interactions dominates the uncertainty about the degree of influence that human activities have on climate. Because clouds form in ascending air currents, whereas cloud droplets nucleate on aerosols that serve as cloud condensation nuclei (CCN), we need accurate measurements of both updrafts and CCN supersaturation ( $S$ ) spectra before we can disentangle aerosol effects on cloud radiative forcing (CRF) from dynamical effects.

## Need for Global Measurements of Cloud Base Updrafts and $CCN(S)$

Tackling the global change problems as identified by the IPCC requires that these quantities be measured on a global scale. However, satellites have not been able to measure updraft speed of the air that forms the clouds or the concentrations of aerosols that are capable of forming cloud drops, which are ingested into the clouds as they grow. Lack of such fundamental quantities has greatly hindered our capability of disentangling the effects of meteorology and anthropogenic aerosol emissions on cloud properties (2). This situation is starting to change with our recently developed methodology to retrieve updrafts at cloud base (3, 4) using the Visible/Infrared Imager Radiometer Suite (VIIRS) instrument onboard the Suomi National Polar-orbiting Partnership (NPP) satellite. This satellite is sun-synchronous, with an overpass time near 13:30 solar time.

Missing such fundamental quantities as  $CCN(S)$  and cloud base updraft  $W_b$  has been preventing us from disentangling the effects of aerosols from atmospheric dynamics (i.e., meteorology). Their absence also has limited our ability to validate the hypothesized impacts of added aerosols on a large range of phenomena, including (i) maintaining full cloud cover in marine stratocumulus, thus incurring a strong cooling effect on the climate system (5); (ii) suppressing precipitation from shallow clouds (6–8); (iii) invigorating the convection in deep tropical clouds (9); (iv) enhancing cloud electrification (10, 11); (v) intensifying severe convective storms to produce more large hail and tornadoes (12); and (vi) decreasing the intensity of tropical cyclones (13). In addition to their intrinsic importance, these aerosol effects could induce radiative effects that change Earth's energy budget in a significant way (1).

Previous satellite-based studies related cloud properties mostly to the aerosol optical depth (AOD) and the Ångström coefficient (14–18). However, AOD as a proxy for CCN is a rather crude tool that is fraught with problems (19) due to a large number of reasons, including (i) aerosol swelling with high relative humidity (20, 21); (ii) uncertainty in solubility and size distribution (18); (iii) lack of a discernible optical signal from small CCN; (iv) cloud contamination (22); (v) AOD not representing aerosol concentrations near cloud base; (vi) cloud obscuration of the aerosols in the boundary layer; (vii) cloud detrainment of aerosols aloft (23, 24) yielding an increase in AOD for deeper and more extensive clouds without corresponding increase in cloud base aerosol concentrations; and (viii) lack of accurate AOD signal for the pristine boundary layer, where accuracy is most critical because clouds respond to the relative change in CCN concentrations, which can be a very small absolute change at very low absolute concentrations (25). These factors often explain a substantial part of the indicated associations of AOD with cloud top properties (18, 26), which has been erroneously ascribed to aerosol effects.

This paper results from the Arthur M. Sackler Colloquium of the National Academy of Sciences, "Improving Our Fundamental Understanding of the Role of Aerosol–Cloud Interactions in the Climate System," held June 23–24, 2015, at the Arnold and Mabel Beckman Center of the National Academies of Sciences and Engineering in Irvine, CA. The complete program and video recordings of most presentations are available on the NAS website at [www.nasonline.org/Aerosol\\_Cloud\\_Interactions](http://www.nasonline.org/Aerosol_Cloud_Interactions).

Author contributions: D.R. designed research; D.R., Y. Zheng, Y. Zhu, Z.L., and M.O.A. performed research; X.Y., G.L., Z.Y., B.F., P.A., H.M.J.B., and U.P. contributed new analytic tools; Y. Zheng, E.H., M.L.P., A.J., C.P., X.Y., Y. Zhu, D.G., and T.G. analyzed data; D.R., U.P., and M.O.A. wrote the paper; M.L.P., A.J., and C.P. provided CCN data; and P.A. and H.M.J.B. facilitated the measurements of CCN in Brazil.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. Email: [daniel.rosenfeld@huji.ac.il](mailto:daniel.rosenfeld@huji.ac.il).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1514044113/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1514044113/-DCSupplemental).

Aerosol optical properties are useful for measuring aerosol type and particle size, which can be identified by active sensor backscatter intensity, color ratio, and polarimetry, or by passive multi-angle intensity measurements even without polarimetry. Adding polarimetry to passive, multiangle imaging should improve the precision and range of conditions under which particle size, shape, and refractive indices can be retrieved. However, this still leaves most of the issues unresolved, especially issues *iii* and *v–viii*, as listed above. To overcome this conundrum, a complete shift in approach is needed. Instead of addressing the limited information content in the optical signal of the aerosols, we extract  $CCN(S)$  by using clouds as an analog for CCN counter (CCNC) chambers.

The structure of this paper is as follows: This section provides the importance and motivation for retrieving  $CCN(S)$ . *Methodology* provides a summary of the recent advancements which constitute a critical mass enabling satellite-only retrieval of  $CCN(S)$  and applies it while describing the essence of the methodology. An extensive validation effort is described in *Validation of the Satellite-Retrieved CCN(S)*, and its results are given in *Results*, along with error calculations. The possibilities that open up with the emerging capabilities for coincident satellite retrieval of convective cloud base updrafts and  $CCN(S)$  are discussed in *Applications of Satellite-Retrieved Updrafts*. Finally, the conclusions are given in *Conclusions*.

## Methodology

**Using Clouds as CCN Chambers.** The commonly used CCNCs measure the number concentration of aerosol particles in a sample air stream ( $N_a$ ), which at a given  $S$  can be activated into the same number of cloud droplets at its base ( $N_{db}$ ) (27). Alternatively, retrieving  $N_{db}$  and  $S$  in clouds can provide  $CCN(S)$ . The peak vapor supersaturation at an adiabatic cloud base,  $S$ , is determined by  $CCN(S)$  and  $W_b$ . Therefore, a good approximation of  $S$  can be calculated from the retrieved  $N_{db}$  and  $W_b$  according to

$$S = C(T_b, P_b) W_b^{3/4} N_{db}^{-1/2}, \quad [1]$$

where  $C$  is a coefficient that depends weakly on cloud base temperature ( $T_b$ ) and pressure ( $P_b$ ) (28). This is an analytical expression that was derived based on theoretical considerations. Recently, it has become possible to estimate  $N_{db}$  and  $W_b$  from satellite measurements, thus calculating also  $S$ . This constitutes the ability to calculate  $CCN(S)$  from satellite measurements only. The following subsections describe the methodology of satellite estimation of  $N_{db}$  and  $W_b$ .

**Estimation of Cloud Base Drop Concentrations.** Retrieving  $T_b$ ,  $P_b$ ,  $W_b$ , and  $N_{db}$  became possible with the advent of the Suomi NPP satellite, which was launched in October 2011. The VIIRS onboard this satellite has a moderate spatial resolution of 750 m. The VIIRS has an Imager with a subset of five channels with double resolution of 375 m at 0.64  $\mu\text{m}$ , 0.865  $\mu\text{m}$ , 1.61  $\mu\text{m}$ , 3.74  $\mu\text{m}$ , and 11.45  $\mu\text{m}$ . Although VIIRS Imager 375-m data were not designed for retrieving cloud properties, a methodology was developed for using it to retrieve cloud drop effective radius ( $r_e$ ) and cloud-top temperatures ( $T$ ). The retrieval of  $r_e$  was based on the methodology developed by Rosenfeld and Lensky (29) for the Advanced Very High Resolution Radiometer. It has been applied to VIIRS by Rosenfeld et al. (30). The ability to retrieve cloud properties at a resolution of 375 m is a breakthrough compared with the previous best available resolution of 1 km. This allows microphysical monitoring of cloud properties with unprecedented accuracy and makes it possible to obtain the microstructure of small clouds at the top of the boundary layer (30).

A VIIRS-retrieved  $T-r_e$  relationship, which is obtained from a convective cloud ensemble within an area of  $\sim 30 \times 30$  km (28), serves as the basis for retrieving  $T_b$ ,  $P_b$ , and  $N_{db}$ . This satellite method is based on extensive aircraft measurements of  $T-r_e$  relationships. It was demonstrated that  $r_e$  varies with altitude nearly as in an adiabatic cloud, and therefore adiabatic cloud drop number concentrations ( $N_{da}$ ) can be calculated at different altitudes in the cloud using the calculated adiabatic water content  $LWC_a$  and  $r_{ea}$ , which is assumed to vary adiabatically with altitude, thus marked as  $r_{ea}$  (31). Then,  $N_{da}$  approximates  $N_{db}$ , because the cloud can be assumed to be adiabatic at its base. The retrieved  $r_e$  is assumed to be  $r_{ea}$  based on the assumption that the measured  $r_e$  is adiabatic, which is the case for clouds with extreme inhomogeneous mixing and with all cloud drops nucleated at their base.

Deviations from the extreme inhomogeneous assumption lead to a reduction of the aircraft-based calculation of  $N_{da}$  by an average factor of 1.3 with respect to the value calculated under this assumption (31). The cloud base drop concentration is approximated by the adiabatic cloud drop concentration as calculated by Eq. 2 (32),

$$N_{da} = \alpha^3 LWC_a / r_{ea}^3, \quad [2]$$


$$\alpha = 62.03 r_e / r_v, \quad [3]$$

where  $r_v$  is the cloud drop mean volume radius, as calculated by equally distributing cloud water content between the cloud droplets. We used the value of  $r_e = 1.08 r_v$  (31). The adiabatic water is obtained from the VIIRS-measured  $T_b$ , which is simply the warmest cloudy pixel, based on a specially developed cloud mask (33). The  $LWC_a$  is calculated based on an adiabatic parcel that rises from cloud base at  $T_b$  and  $P_b$  to the isotherm  $T$ , where multiple pairs of  $T$  and  $r_{ea}$  are retrieved for different cloudy pixels of the same cloud cluster at different heights above cloud base. Here,  $P_b$  is obtained from the pressure at the isotherm of satellite-retrieved cloud base height ( $H_b$ ), which was computed from the European Center for Medium-range Weather Forecasting reanalysis data.  $H_b$  was calculated as the difference between reanalysis surface air temperature and  $T_b$  divided by the dry adiabatic lapse rate.  $T_b$  was validated at a root-mean-square (RMS) error of 1.1 K, as shown in Fig. 1 (33).  $H_b$  and  $T_b$  were calculated for conditions of convective clouds that developed from well-mixed boundary layer at the early afternoon satellite overpass time (33), before the peak of the convective rain and the resultant cooling and moistening of the boundary layer by evaporating precipitation.

**Estimation of Cloud Base Updrafts.** Until now, only lidar and radar measurements of  $W_b$  were used. This is expanded here to satellite-retrieved  $W_b$ . According to Eq. 1, knowing  $W_b$  and  $N_{db}$  at cloud base yields  $S$ . Then,  $N_{db}$  is numerically identical to  $CCN(S)$ . Rosenfeld et al. (32) used this method to retrieve  $CCN(S)$  over the Atmospheric Radiation Measurement (ARM) site of the Southern Great Plains (SGP), using  $N_{db}$  retrieved from a satellite and  $W_b$  measured by ARM's vertically pointing  $K_a$ -band radar. The  $W_b$  was calculated from all full Doppler statistics during a 2-h window centered at the satellite overpass time, where the  $W_b$  of each point in time was weighted by  $W_b$  itself, thus representing its relative contribution to building the cloud volume. More specifically, equation 5 in Rosenfeld et al. (32) (replicated as Eq. 4 here) shows that the radar or lidar updraft  $W$  was constructed from all of the  $N$  realizations  $W_i$  of single data points within the time window as follows:

$$W = \frac{\sum N_i W_i^2}{\sum N_i W_i} \Bigg|_{W_i > 0} \quad [4]$$

According to Eq. 4,  $W$  is the cloud volume-weighted updraft. Good agreement was achieved by Rosenfeld et al. (32) between  $CCN(S)$  as



**Fig. 1.** The relationship between satellite-measured cloud base temperature and validation measurements by a combination of a ceilometer and soundings at the Department of Energy (DOE)/Atmospheric System Research (ASR) sites on the SGP in Oklahoma. Reproduced from ref. 33.

constructed by satellite-retrieved  $N_b$  and radar-retrieved  $W_b$  with the SGP ground base-measured  $CCN(S)$ , but the number of cases with useful clouds and data were rather small and served mainly to verify the methodology. The need for ground-based measurements of  $W_b$  limited severely the occasions where  $CCN$  could be retrieved to sites where cloud Doppler lidars or radars measurements are available. The present study is the first one, to our knowledge, to retrieve  $CCN(S)$  from satellite estimates of both  $N_{db}$  and  $W_b$ , thus becoming potentially very widely applicable, despite some limitations in the retrievals of  $N_{db}$  and  $W_b$ .

Retrieval of  $CCN$  solely from satellite data requires  $W_b$  to be retrieved from satellite. This was done by using satellite-retrieved components of the energy that propels the convection (3). Subsequently, Zheng and Rosenfeld (4) showed that  $W_b$  can be simply calculated by

$$W_b = AH_b, \quad [5]$$

where  $W_b$  is cloud base updraft in meters per second,  $A$  is a coefficient ( $0.0009 \text{ s}^{-1}$ ) obtained in a previous study (4), and  $H_b$  is the cloud base height above the ground in meters, which is determined by the difference between the surface air and cloud base temperatures, as explained at the end of *Estimation of Cloud Base Drop Concentrations*. This relationship was developed based on synchronous satellite and lidar measurements from the ARM SGP site and at the ARM Mobile Facility onboard a ship on a line between Los Angeles and Honolulu [Marine ARM GPCI Investigations of Clouds (MAGIC)]. The satellite-retrieved  $W_b$  was validated against the Doppler measurements, resulting in an RMS error of  $0.41 \text{ m s}^{-1}$  and a mean absolute percentage error (MAPE) of 24% and 21% by Zheng et al. (3) and Zheng and Rosenfeld (4), respectively. When forcing the relationships through zero (Eq. 5 and Fig. 2), the error becomes 27%. These results are consistent with the physical considerations of Williams and Stanfill (10). This means that the methodology is very likely to be universally applicable to boundary layer convective clouds.

Table 1 summarizes the methodology. It shows the satellite measurements, their combination with reanalysis data, and their propagation into the eventual  $W_b$  and  $CCN(S)$ , and the associated errors.

### Validation of the Satellite-Retrieved $CCN(S)$

Cloud base  $S$  was obtained from Eq. 1, with  $N_{db}$  calculated by Eq. 2 and  $W_b$  calculated using Eq. 5. The calculated  $N_{db}$  is by definition equal to  $CCN(S)$  at cloud base. To compare with surface-based measurements, the concentration is corrected for the difference between air density at cloud base and at the ground, and then validated against the  $CCN(S)$  as measured by the ground-based instrument. This assumes that the thermals bring the surface air to cloud base without much change in the mixing ratio and properties of aerosol particles. This is a widely accepted assumption for vapor mixing ratio at thermally driven cloud bases in a well-mixed


boundary layer, where the lifting condensation level is usually very similar to the actual cloud base height.

An initial comparison of the satellite-retrieved  $CCN$  to the SGP instrumental validation data (assuming no error in the instrument measured  $CCN$ ) showed a slope of 0.74 for the regression line. A retrieval bias could be caused by a large number of factors, which are quantified in *Error Analysis*, but the largest potential source of error is inaccuracy in  $r_c$ . The observed 26% underestimate in  $CCN$  could have been caused by a 10% systematic overestimate in the retrieved  $r_c$ . This is quite probable, because MODIS-retrieved  $r_c$  was found to be larger by 10–15% than aircraft in situ measurements (34–36). An underestimate of satellite- versus surface-measured  $CCN$  can be also caused by a systematic decrease of  $CCN$  number concentration ( $N_{CCN}$ ) between the surface and cloud base heights. This bias has to be corrected before calculating  $S$  by Eq. 1, because, otherwise,  $S$  would be overestimated. To stay on the conservative side, we applied only half of the bias correction and used here a reduction factor of 1.15 instead of 1.3, as proposed by Freud et al. (31), and applied it to all of the validation sites.

Validation cases were selected over the sites of the ARM sites of the SGP in Oklahoma, at Manacapuru near Manaus in the Amazon, and over the northeastern Pacific onboard the MAGIC ship. In addition,  $CCN$  measurements were obtained from the Amazon Tall Tower Observatory (ATTO) site 150 km to the northeast of Manaus (37). Data were obtained from the start of availability of VIIRS data in 2012 until early 2015. The case selection criteria were as follows: (i) Satellite overpass has to occur at a zenith angle between  $0^\circ$  and  $45^\circ$  to the east of the ground track, which is the sunny side of the clouds. For a specific location, these satellite views occur once or twice every 6 d. (ii) Convective clouds must occur and possess a vertical development that spans at least 6 K of cloud temperature from base to top, limiting to clouds with thickness  $>1$  km. (iii) The clouds must not precipitate significantly (i.e., without a radar or lidar detectable rain shaft that reaches the ground). The precipitation causes cold pools that disconnect the continuity of the air between the surface and the cloud base. (iv) Cloud elements with indicated  $r_c > 18 \mu\text{m}$  are rejected automatically from the analysis that is likely to rain/drizzle heavily. (v) No obscuration from high clouds is allowed. An automatic detection of semitransparent clouds screens them from the selected area for analysis. (vi) Ground-based  $CCN$  data must be available.

The availability of  $CCN$  data from the ARM program at all of its three sites was severely limited due to data quality issues. Insufficient available time for stabilization of temperatures at low  $S$  caused the  $CCN$  readings at  $S \leq 0.25\%$  to be grossly underestimated or zero, and therefore they could not be used. The points with  $S > 0.25\%$  were fit with a second-order polynomial that was forced through the origin, because  $CCN$  must be zero for  $S = 0$ . By extrapolation with this polynomial, we could extend the use of the data down to  $S = 0.2\%$ . Cases with cloud base  $S < 0.2\%$  were rejected. The operation of the ARM CCNCs was changed after August 2014 to allow sufficient time for stabilization at low  $S$ . This correction was applied to Manacapuru only by April 2015, however. These limitations did not apply to ATTO, and valid data from this site were available from May 2014 until January 2015.

The results are shown in Fig. 3. Each point in the figure represents one satellite overpass over one ground-based  $CCN$ . The  $CCN$  data from a time window of  $\pm 1$  h around the overpass are taken to include several  $CCN(S)$  spectra at all measured supersaturations. Because of the much slower scanning rate of  $S$  at ATTO, a larger time window of  $\pm 1.5$  h was taken there to include at least one full spectrum of  $CCN(S)$ . The satellite analyzes clouds over an area of about  $30 \times 30$  km around the ground measurement site, with some adjustments to incorporate the convective clouds in the vicinity. The satellite-retrieved  $CCN$  and  $S$  are compared with the instrument measurements as follows:



**Fig. 2.** Relationship between satellite-estimated  $W_b = 0.0009H_b$  and directly measured  $W_b$  by Doppler lidars at the SGP (red dots), Green Ocean Amazon (GOAmazon) (green), and MAGIC (blue). After Zheng and Rosenfeld (4). The satellite-retrieved  $T_b$  was converted to height using the European Centre for Medium-Range Weather Forecasts reanalysis data. RMSE is the estimation root mean square error.

**Table 1. Propagation of the calculations from the satellite retrievals to the resultant  $CCN(S)$** 

Symbol	Parameter and unit	Source of calculation	Error
$r_e$	cloud drop effective radius, $\mu\text{m}$	satellite retrieval	8%
$T$	cloud surface temperature, $^{\circ}\text{C}$	satellite retrieval	0.2 $^{\circ}\text{C}$
$T_b$	cloud base temperature, $^{\circ}\text{C}$	satellite retrieval	1.1 $^{\circ}\text{C}$
$P_b$	cloud base pressure, hPa	$T_b$ + reanalysis	15 hPa
$r_v$	drop mean volume radius, $\mu\text{m}$	$r_e$ (Eq. 3)	8%
$LWC_a$	cloud adiabatic water, $\text{g}\cdot\text{kg}^{-1}$	$T + T_b + P_b$ (parcel)	15%
$N_d$	cloud base drop concentrations, $\text{cm}^{-3}$	$r_v(T) + LWC_a(T)$ (Eq. 2)	30%
$H_b$	cloud base height above surface, m	$T_b$ + reanalysis	150 m
$W_b$	cloud base updraft, $\text{m}\cdot\text{s}^{-1}$	$H_b$ (Eq. 5)	27%
$S$	cloud base max supersaturation, %	$T_b, P_b, W_b, N_d$ (Eq. 1)	25% of $S$ in percent
$N_{CCN}(S)$	CCN at cloud base, $\text{cm}^{-3}$	$N_d, S$ by definition	30%


(i) A scatter plot of the individual ground-based measurements of CCN concentrations ( $N_{CCN}$ ) is plotted as a function of  $S$ . (ii) A second-order polynomial curve is fit to the points. The function is forced through the origin, because zero  $S$  must correspond to zero  $N_{CCN}$ . (iii) The  $N_{CCN}$  is taken from the polynomial fit at the same  $S$  that is retrieved from satellite at cloud base. The  $\pm 95\%$  confidence interval of  $N_{CCN}$  at the value of satellite-retrieved  $S$  is calculated. (iv) The satellite-retrieved  $N_{CCN}$  is the satellite-retrieved  $N_{db}$ , corrected for the air density difference between cloud base and the surface.

### Results

Fig. 3 shows the relationships between the satellite retrievals of  $N_{CCN}$  and  $S$  at cloud base, and the ground-based measurements

of  $N_{CCN}$  at the same  $S$ . There are several points worth noting. (i) The figure covers a large dynamic range of  $S$  for both low and high values  $N_{CCN}$ . (ii) The value of  $R^2 = 0.76$  shows that the fit explains more than 3/4 of the variability between the satellite and ground-based measurements of  $CCN(S)$ . (iii) There is a systematic underestimate bias of 14% in the satellite-retrieved CCN. It follows that the estimation errors decrease almost linearly with smaller  $N_{CCN}$ . (iv) The variation of the satellite with respect to the ground-based measurements is within 20–25% of the ground-based measurements. This includes the 14% bias error. (v) The SD of the fit is similar to the expected magnitude from the error sources of the satellite uncertainties in  $W_b, T_b,$  and  $r_e$ .

The methodology was converted into a procedure that can be applied to any specified rectangle in the VIIRS imagery, which



**Fig. 3.** The relationship between satellite-retrieved  $N_{CCN}$  and  $S$  at cloud base, and the ground-based instrument measurements of  $N_{CCN}$  at the same  $S$ . The slope and intercept of the best fit line are given in the key by  $m$  and  $b$ , respectively. The validation data are collected from the DOE/ASR sites on the SGP in Oklahoma and GOAmazon near Manaus, and over the northeast Pacific (MAGIC). In addition, data are obtained from the ATTO. The location is denoted by the marker shape, and  $S$  is shown by the color.


contains surface thermally driven convective clouds, and provides as output the following parameters:  $T_b$ ,  $P_b$ ,  $H_b$ ,  $W_b$ ,  $N_{db}$ , and  $S$ . The value of  $N_{db}$  is equal to the CCN concentrations at the retrieved  $S$  at cloud base, and this value of  $CCN(S)$  is also an output parameter. As an illustrative example, this procedure was applied to a regular grid of  $75 \times 75$  VIIRS Imager pixels ( $28 \times 28$  km at nadir) over the region of Houston during conditions of onshore flow of a tropical marine air mass. The results are shown in Fig. 4. The salient features are as follows: (i) There are very low CCN concentrations over the ocean. (ii) There is only a modest increase in CCN over the rural areas inland. (iii) The CCN concentrations more than triple over and downwind of the urban area compared with the cross-wind areas. (iv)  $S$  decreases over the urban area to less than half of the values over the rural areas. Therefore, CCN for the same  $S$  is enhanced by a factor much larger than 3. (v) The indicated CCN concentrations are similar in adjacent areas with similar conditions, indicating the robustness of the methodology.

### Applications of Satellite-Retrieved Updrafts and $CCN(S)$ to Reduced Climate Uncertainties

Here we showed the feasibility to retrieve  $CCN(S)$  from a single satellite passive sensor using clouds as CCN chambers, under certain conditions. There are still many challenges to overcome before it will be possible to do so for most cloud types. This requires the development of new satellite capabilities that will be

able to provide more direct measurements of updraft speeds, such as measuring vertical motions of cloud elements by tracking their evolution with time. Here we attempt to open a window to the potential applications of such capability, with a few examples.

The sensitivity of cloud properties to  $N_{CCN}$  is logarithmic (38). This means that a small absolute change in  $N_{CCN}$  has a much larger impact during pristine than polluted conditions. Carslaw et al. (25) argued that the main sensitivity to anthropogenic aerosols occurs in areas that had  $N_{CCN}$  of  $35\text{--}65\text{ cm}^{-3}$  during the preindustrial era. Satellite measurements show that an increase of more than  $100\text{ W}\cdot\text{m}^{-2}$  in cloud radiative effect (CRE) can occur when  $N_d$  of marine shallow boundary layer clouds increases from  $35\text{ cm}^{-3}$  to  $65\text{ cm}^{-3}$ , mainly due to increased cloud cover and cloud liquid water path. This is manifested as closing areas of open cellular convection (39). However, the satellite observed  $N_d$  is related to both  $W_b$  and  $CCN(S)$ , as shown by Eq. 1. Therefore, there is a possibility that measurements of the large enhancement of CRE that were associated with increased  $N_d$  could also result from changes in  $W_b$ , which could be caused by changes of meteorology (40). For separating the roles of  $W_b$  and  $N_{CCN}$  in the determination of  $N_{db}$ , both  $W_b$  and  $N_{CCN}$  should be measured. As already discussed in *Need for Global Measurements*, using AOD as a proxy for  $N_{CCN}$  in the marine boundary layer clouds has several shortcomings. Because, among other problems, the correlation between AOD and  $N_{CCN}$  is not very close and because a column property like AOD is not necessarily representative of the CCN



**Fig. 4.** Application of the methodology to the Houston area. The retrieval is done for a regular grid of  $75 \times 75$  375-m VIIRS/Imager pixels ( $\sim 28 \times 28$  km at nadir). The numbers in each area are: top, CCN (per cubic centimeter); middle,  $S$  (percent); and bottom, cloud base temperature (degrees Celsius). Unstable clean tropical air mass flows northward (upward in the image) from the Gulf of Mexico. The Houston urban effect is clearly visible by more than tripled CCN concentrations over Houston and the reduction of  $S$  to less than half. This represents an even larger factor in enhancing CCN for the same  $S$ . A smaller effect is seen over the urban and industrial areas to the east of Houston. The color composite is red, green, and blue for the visible reflectance,  $3.7\text{-}\mu\text{m}$  solar reflectance, and thermal temperature, respectively, as in Rosenfeld et al. (30). The Houston bay and beltways are marked by white lines.

concentrations that affect growing clouds, the AOD approach allows only an order-of-magnitude estimate of  $N_{CCN}$ . On the other hand, combining  $W_b$  with  $N_{db}$  can provide  $CCN(S)$  with an uncertainty that can be quantified and is far better than the AOD approach. Having both  $W_b$  and  $CCN(S)$  will allow disentangling of the roles of these two factors in determining  $N_d$  and in the attribution of the related changes in CRE to aerosols.

Having satellite retrievals of both  $W_b$  and  $N_{CCN}$  will allow disentangling of their respective roles on determining  $N_d$  and the related precipitation-forming processes, rainfall amounts, and distribution of vertical latent heating.  $CCN(S)$  ingested by deep convective clouds can be estimated by using adjacent shallower nonprecipitating convective clouds in their upwind side. Adding CCN to deep convective clouds can invigorate them and incur more extensive anvils and respective positive radiative forcing (41–43). This can be quantified observationally using long-term surface aerosol, cloud, and meteorological measurements made at a single location in the SGP (41, 42), and also using global A-Train satellite products (44). These estimates of CRF (the change in CRE due to anthropogenic causes) are associated with aerosol-induced changes in cloud properties that do not differentiate the respective roles of aerosol and dynamics or meteorology but their joint effects.

Having global coverage of  $CCN(S)$  where we need them most—in conjunction with the clouds that ingest them—will provide input for regional and global simulations. The coincident retrieved cloud properties will constrain these models and provide us with realistic assessments of the CRE. The retrieved  $CCN(S)$  can be used for constraining aerosol production and transport models. This will allow separating the aerosols into natural and anthropogenic components more accurately. The application of such classified  $CCN(S)$  will facilitate calculating the anthropogenic aerosol-induced CRF, which will constitute a major reduction of the uncertainty in anthropogenic climate forcing. Developing methodologies for retrieving updrafts at clouds with elevated bases will further enhance these possibilities.

## Conclusions

The feasibility of estimating  $CCN(S)$  and  $W_b$  of boundary layer clouds from the Suomi NPP polar-orbiting operational weather satellite was demonstrated with an accuracy of  $\pm 25\%$  to  $\pm 30\%$ , which is limited mostly by the accuracy in the retrieval of  $r_e$ . The validation was done in Oklahoma, the Amazon Basin, and the northeast Pacific Ocean. Our methodology is presently limited to boundary layer convective clouds of at least 1 km depth, which are not obscured by upper layer clouds, including semitransparent cirrus. This might limit its application in some regions of the world. Moreover, the limitation for small solar backscattering angles of  $<25^\circ$  restricts the satellite coverage to 1/4 of the satellite swath width, or a view once every 4 d, on average. On the other hand, even for a regional coverage, it would be much more valuable to study the process of aerosol–cloud interactions rather than using single-point data as provided by ground-based observations.

A major advantage of using clouds as analog for CCN chambers relative to relying on the optical signal of the aerosols themselves is the fact that the optical signals (e.g., AOD and Ångström coefficient) vanish at very small aerosol concentrations, which is exactly where the relative changes in CCN concentrations matter most, or, in other words, where very small absolute changes in concentrations have very large impacts on clouds (16, 25). This is where the traditional remote sensing methods of aerosols break down, whereas the applicability of using clouds as CCN chambers remains intact, as evident by the lower left corner of Fig. 3. This has particular importance in the context of the quest for the significance of changes from the preindustrial era to the present background aerosols (25).

The retrieval of both  $CCN(S)$  and  $W_b$  allows, for the first time to our knowledge, disentangling of the roles of updrafts and

CCN on cloud microphysical, precipitation, and radiative properties over climatically meaningful areas. Previously, the inability to separate these factors has been a major impediment to our ability to quantify the aerosol/cloud-mediated effects on Earth's energy budget, thus keeping high the uncertainty of this effect (1). Application of the new capabilities offered by our methodology is expected to allow a breakthrough in quantifying these effects and to substantially reduce the uncertainty in anthropogenic aerosol climate forcing, at least for boundary layer convective clouds.

## Error Analysis

A direct comparison of the satellite- to ground-based CCN, assuming no errors in the CCN measurements, shows a correlation coefficient of 0.88 and a slope of 0.9 (i.e., underestimate of 10%). The MAPE is  $\pm 30\%$ . However, both satellite retrievals and CCN measurements are subject to errors. Therefore, a bivariate regression has to be used for fitting two parameters with associated errors for both (45). The associated error for the satellite-retrieved CCN for a given  $S$  was taken as  $\pm 30\%$ . The CCN instrument errors were taken as the  $\pm 95\%$  confidence interval (i.e.,  $\pm 2$  SDs) of  $N_{CCN}$  for the individual cases, as described at the end of *Estimation of Cloud Base Drop Concentrations*. Both sets of errors are shown as error bars in Fig. 3.

The largest sensitivity is to errors in  $r_e$ , because, according to Eq. 2, the error in  $N_a$  is the cube of the error in  $r_e$ . The accuracy of MODIS-retrieved  $r_e$  is best when the 3.7- $\mu\text{m}$  waveband is used (MODIS  $r_e$  is also available for 2.1- $\mu\text{m}$  and 1.6- $\mu\text{m}$  wavebands) in nondrizzling clouds; under these conditions, it showed the best agreement with aircraft measurements, with an uncertainty of 1  $\mu\text{m}$  (46). The 3.7- $\mu\text{m}$ -based  $r_e$  is also minimally affected by cloud inhomogeneities (47) because this band absorbs solar radiation much more strongly (48, 49). The VIIRS footprint area, which is sevenfold smaller than that of MODIS, further reduces the possibility of errors caused by cloud inhomogeneities. Our implementation to VIIRS is even more accurate than MODIS in the best of circumstances, because we use only pixels with visible reflectance  $>0.4$  at backscattering angles (satellite zenith angle of 0–50 degrees). To avoid significant distortion of  $r_e$  by coalescence, we avoided heavily precipitating clouds at their tops ( $r_e > 18 \mu\text{m}$ ). MODIS  $r_e$  is larger than aircraft in situ measurements by 10–15% (34–36). This is probably not a problem for retrieved  $r_e$  based on the VIIRS Imager (30), because it is lower by a similar amount with respect to MODIS  $r_e$ . The retrieval uncertainty of  $r_e$  itself is roughly  $\pm 10\%$  (36). This translates to uncertainty of a factor of  $\pm 33\%$  in  $N_a$ . This error alone is larger than the measured validation error of  $\pm 30\%$  when assuming no errors in the ground-measured CCN, which includes many other error sources, as described next. This might serve as an indication that the error in the retrieved  $r_e$  from VIIRS is smaller than for MODIS, probably due to the much finer resolution.

The MAPE in cloud base temperature of  $\pm 1.1^\circ\text{C}$  propagates to a 5% error in  $N_a$  due to changing  $C(T_b, P_b)$  in Eq. 1. The error in  $W_b$  (Fig. 2) can be propagated to an error in  $N_a$  according to Twomey's approximation of

$$N_a = CCN(S = 1\%)^{2/(k+2)} W_b^{3k/(2k+4)}, \quad [6]$$

where  $k$  is the slope of the  $CCN(S)$  spectrum on a log–log scale (50). Accordingly, a  $W_b$  MAPE of  $\pm 27\%$  propagates to an error in  $N_a$  of only 7–13% for  $k = 0.5$  and 1, respectively. The overall combined error is  $\pm 36\%$ , as obtained by the calculation  $(0.33^2 + 0.05^2 + 0.13^2)^{0.5} = 0.36$ . This overall calculated error of  $\pm 35\%$ , even before adding the CCN instrument uncertainty, is larger than the measured validation error of  $\pm 30\%$  when assuming no errors in the ground-measured CCN. This discrepancy could be explained, for example, by reducing the  $r_e$  error from 10% to 8%.

**ACKNOWLEDGMENTS.** This study was made possible by merging the efforts of many researchers around the world, with a respective long list of acknowledgments. The study was supported by the US Department of Energy Office of Science Atmospheric System Research program (Grant DE-SC0006787), the State Basic Research Development Program of China (2013CB955804), and National Oceanographic and Atmospheric Administration Joint Polar Satellite System Proving Ground and Risk Reduction program. This study is also partially supported by project BACCHUS

- Boucher O, et al. (2013) Clouds and aerosols. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge Univ Press, New York), pp 571–657.
- Rosenfeld D, Sherwood S, Wood R, Donner L (2014) Atmospheric science. Climate effects of aerosol-cloud interactions. *Science* 343(6169):379–380.
- Zheng Y, Rosenfeld D, Li Z (2015) Satellite inference of thermals and cloud base updraft speeds based on retrieved surface and cloud base temperatures. *J Atmos Sci* 72(6):2411–2428.
- Zheng Y, Rosenfeld D (2015) Linear relation between convective cloud base height and updrafts and application to satellite retrievals. *Geophys Res Lett* 42(15):6485–6491.
- Goren T, Rosenfeld D (2012) Satellite observations of ship emission induced transitions from broken to closed cell marine stratocumulus over large areas. *J Geophys Res* 117(D17):D17206.
- Rosenfeld D (1999) TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall. *Geophys Res Lett* 26(20):3105–3108.
- Rosenfeld D (2000) Suppression of rain and snow by urban and industrial air pollution. *Science* 287(5459):1793–1796.
- Andreae MO, et al. (2004) Smoking rain clouds over the Amazon. *Science* 303(5662):1337–1342.
- Rosenfeld D, et al. (2008) Flood or drought: How do aerosols affect precipitation? *Science* 321(5894):1309–1313.
- Williams E, Stanfill S (2002) The physical origin of the land–ocean contrast in lightning activity. *C R Phys* 3(10):1277–1292.
- Bell TL, Rosenfeld D, Kim KM (2009) Weekly cycle of lightning: Evidence of storm invigoration by pollution. *Geophys Res Lett* 36(23):L23805.
- Rosenfeld D, Bell TL (2011) Why do tornados and hailstorms rest on weekends? *J Geophys Res* 116(D20):D20211.
- Rosenfeld D, et al. (2012) Aerosol effects on microstructure and intensity of tropical cyclones. *Bull Am Meteorol Soc* 93(7):987–1001.
- Koren I, Kaufman YJ, Rosenfeld D, Remer LA, Rudich Y (2005) Aerosol invigoration and restructuring of Atlantic convective clouds. *Geophys Res Lett* 32(14):L14828.
- Koren I, et al. (2012) Aerosol-induced intensification of rain from the tropics to the mid-latitudes. *Nat Geosci* 5(2):118–122.
- Koren I, Dagan G, Altaratz O (2014) From aerosol-limited to invigoration of warm convective clouds. *Science* 344(6188):1143–1146.
- Niu F, Li Z (2012) Systematic variations of cloud top temperature and precipitation rate with aerosols over the global tropics. *Atmos Chem Phys* 12(18):8491–8498.
- Liu J, Li Z (2014) Estimation of cloud condensation nuclei concentration from aerosol optical quantities: Influential factors and uncertainties. *Atmos Chem Phys* 14(1):471–483.
- Andreae MO (2009) Correlation between cloud condensation nuclei concentration and aerosol optical thickness in remote and polluted regions. *Atmos Chem Phys* 9(2):543–556.
- Kapustin V, et al. (2006) On the determination of a cloud condensation nuclei from satellite: Challenges and possibilities. *J Geophys Res* 111(D4):D04202.
- Boucher O, Quaas J (2013) Water vapour affects both rain and aerosol optical depth. *Nat Geosci* 6(1):4–5.
- Koren I, Oreopoulos L, Feingold G, Remer L, Altaratz O (2008) How small is a small cloud? *Atmos Chem Phys* 8(14):3855–3864.
- Chakraborty S, Fu R, Wright JS, Massie ST (2015) Relationships between convective structure and transport of aerosols to the upper troposphere deduced from satellite observations. *J Geophys Res* 120(13):6515–6536.
- Zhu Y, Rosenfeld D, Yu X, Li Z (2015) Separating aerosol microphysical effects and satellite measurement artifacts of the relationships between warm rain onset height and aerosol optical depth. *J Geophys Res*, 120(15):7726–7736.
- Carslaw KS, et al. (2013) Large contribution of natural aerosols to uncertainty in indirect forcing. *Nature* 503(7474):67–71.
- Gryspeerd E, Stier P, Grandey B (2014) Cloud fraction mediates the aerosol optical depth-cloud top height relationship. *Geophys Res Lett* 41(10):3622–3627.
- European Commission FP7-603445. Measurements at Amazon Tall Tower Observatory are partially supported by FAPESP (Sao Paulo Research Foundation) Projects 13/50510-5 and 13/05014-0, by the Max Planck Society, and by the German Federal Ministry of Education and Research. We acknowledge the support from Instituto Nacional de Pesquisas da Amazonia and the LBA (The Large-Scale Biosphere-Atmosphere Experiment in Amazonia) central office for ATTO and GOAmazon (Green Ocean Amazon) logistical support and operation.
- Roberts G, Nees A (2005) A continuous-flow streamwise thermal-gradient CCN chamber for atmospheric measurements. *Aerosol Sci Technol* 39(3):206–221.
- Pinsky M, Khain A, Mazin I, Korolev A (2012) Analytical estimation of droplet concentration at cloud base. *J Geophys Res* 117(D18):D18211.
- Rosenfeld D, Lensky IM (1998) Satellite-based insights into precipitation formation processes in continental and maritime convective clouds. *Bull Am Meteorol Soc* 79(11):2457–2476.
- Rosenfeld D, et al. (2014) High-resolution (375 m) cloud microstructure as seen from the NPP/VIIRS satellite imager. *Atmos Chem Phys* 14(5):2479–2496.
- Freud E, Rosenfeld D, Kulkarni J (2011) Resolving both entrainment-mixing and number of activated CCN in deep convective clouds. *Atmos Chem Phys* 11(24):12887–12900.
- Rosenfeld D, Fischman B, Zheng Y, Goren T, Giguzin D (2014) Combined satellite and radar retrievals of drop concentration and CCN at convective cloud base. *Geophys Res Lett* 41(9):3259–3265.
- Zhu Y, et al. (2014) Satellite retrieval of convective cloud base temperature based on the NPP/VIIRS Imager. *Geophys Res Lett* 41(4):1308–1313.
- Painemal D, Zuidema P (2011) Assessment of MODIS cloud effective radius and optical thickness retrievals over the Southeast Pacific with VOCALS-REx in situ measurements. *J Geophys Res* 116(D24):D24206.
- Min Q, et al. (2012) Comparison of MODIS cloud microphysical properties with in-situ measurements over the Southeast Pacific. *Atmos Chem Phys* 12(23):11261–11273.
- Noble SR, Hudson JG (2015) MODIS comparisons with northeastern Pacific in situ stratocumulus microphysics. *J Geophys Res* 120(16):8332–8344.
- Andreae M, et al. (2015) The Amazon Tall Tower Observatory (ATTO): overview of pilot measurements on ecosystem ecology, meteorology, trace gases, and aerosols. *Atmos Chem Phys* 15(18):10723–10776.
- Koren I, Martins JV, Remer LA, Afargan H (2008) Smoke invigoration versus inhibition of clouds over the Amazon. *Science* 321(5891):946–949.
- Goren T, Rosenfeld D (2014) Decomposing aerosol cloud radiative effects into cloud cover, liquid water path and Twomey components in marine stratocumulus. *Atmos Res* 138:378–393.
- George RC, Wood R (2010) Subseasonal variability of low cloud radiative properties over the southeast Pacific Ocean. *Atmos Chem Phys* 10(8):4047–4063.
- Li Z, et al. (2011) Long-term impacts of aerosols on the vertical development of clouds and precipitation. *Nat Geosci* 4(12):888–894.
- Yan H, Li Z, Huang J, Cribb M, Liu J (2014) Long-term aerosol-mediated changes in cloud radiative forcing of deep clouds at the top and bottom of the atmosphere over the Southern Great Plains. *Atmos Chem Phys* 14(14):7113–7124.
- Fan J, et al. (2013) Microphysical effects determine macrophysical response for aerosol impacts on deep convective clouds. *Proc Natl Acad Sci USA* 110(48):E4581–E4590.
- Peng J, Li Z, Zhang H, Liu J, Cribb M (2015) Systematic changes in cloud radiative forcing with aerosol loading for deep clouds in the tropics. *J Atmos Sci* 73(1):231–249.
- Cantrell C (2008) Technical Note: Review of methods for linear least-squares fitting of data and application to atmospheric chemistry problems. *Atmos Chem Phys* 8(17):5477–5487.
- King N, Bower K, Crosier J, Crawford I (2013) Evaluating MODIS cloud retrievals with in situ observations from VOCALS-REx. *Atmos Chem Phys* 13(1):191–209.
- Zhang Z, Platnick S (2011) An assessment of differences between cloud effective particle radius retrievals for marine water clouds from three MODIS spectral bands. *J Geophys Res Atmos* 116(D20):D20215.
- Rosenfeld D, Cattani E, Melani S, Levizzani V (2004) Considerations on daylight operation of 1.6-versus 3.7- $\mu$ m channel on NOAA and METOP satellites. *Bull Am Meteorol Soc* 85(6):873–881.
- Grosvenor D, Wood R (2014) The effect of solar zenith angle on MODIS cloud optical and microphysical retrievals within marine liquid water clouds. *Atmos Chem Phys* 14(14):7291–7321.
- Twomey S (1959) The nuclei of natural cloud formation part II: The supersaturation in natural clouds and the variation of cloud droplet concentration. *Geofis Pura Appl* 43(1):243–249.